

FIGURE 1








FIGURE 2A




FIGURE 2B





FIGURE 3




[illegible]

FIGURE 4B





FIGURE 5




Activity of F1-102. *Column A.* Ramos cells were incubated overnight with 10 ug of Human IgG (filled histogram) or with 10 ug of the purified human antibody indicated to the left of the histogram (lined histogram). *Column B.* Ramos cells were incubated overnight in the presence of CD40L and 10 ug of Human IgG (filled histogram) or with 10 ug of the purified human antibody indicated to the left of the histograms (lined histogram). All cells were stained with PE-labeled anti-CD95. The indicated antibody was purified from ascites.




Activity of F4-465 and F5-152. Column A. Ramos cells were incubated overnight with 10 ug of Human IgG (filled histogram) or with 10 ug of the purified human antibody indicated to the left of the histograms (lined histogram). *Column B.* Ramos cells were incubated overnight in the presence of CD40L and 10 ug of Human IgG (filled histogram) or with 10 ug of the purified human antibody indicated to the left of the histograms (lined histogram). All cells were stained with PE-labeled anti-CD95. The indicated antibodies were purified from ascites.

FIGURE 6



F2-103 (K1H1)





B



F5-157 (K3H3)



F5-77 (K1H1)



Agonistic activity of F2-103, F5-157 and F5-77. Column A., Ramos cells were incubated overnight with 10 ug of Human IgG (filled histogram) or with 10 ug of the purified human antibody indicated to the left of the histograms (lined histogram). Column B, Ramos cells were incubated overnight in the presence of CD40L and 10 ug of Human IgG (filled histogram) or with 10 ug of the purified human antibody indicated to the left of the histograms (lined histogram). All cells were stained with PE-labeled anti-CD95. The indicated antibodies were expressed in transiently transfected Cos cells and Ig was purified from the conditioned media.

FIGURE 7